






















# The Theoretical Description for VO(OH)-Assisted Electrochemical Determination for the Cathodic Removal of Environmentally Unfriendly Artificial Sweetener Sucralose and the Chemical Warfare Agent Chloropicrin

Volodymyr V. Tkach <sup>1,\*</sup>, Tetiana V. Morozova <sup>2</sup>, Nataliia M. Storoshchuk <sup>1</sup>, Bogdan D. Storoshchuk <sup>1</sup>, Sílvio C. de Oliveira <sup>3</sup>, Lucinda Vaz dos Reis <sup>4</sup>, Yana G. Ivanushko <sup>5</sup>, Bohdana Yu. Banul <sup>5</sup>, Tetiana V. Honchar <sup>5</sup>, Jarem R. Garcia <sup>6</sup>, Viktoriia V. Paientko <sup>7</sup>, Petro I. Yagodynets <sup>1,\*</sup>, Zholt O. Kormosh <sup>8,\*</sup>, Karina V. Palamarek <sup>9</sup>, Valentyna A. Chychun <sup>9</sup>, Konon L. Bagrii <sup>9</sup>, José Inácio Ferrão da Paiva Martins <sup>10</sup>, Inesa M. Khmeliar <sup>11</sup>, Lesya O. Kushnir <sup>11</sup>, Rostyslav O. Sabadyshyn <sup>11</sup>, Dmytro L. Lysytsia <sup>11</sup>, Riboba G. Sharipova <sup>12</sup>

<sup>1</sup> Chernivtsi National University, 58000, Kotsyubyns'ky Str. 2, Chernivtsi, Ukraine

<sup>2</sup> National Transport University, 02000, Omelianovych-Pavlenko Str. 1, Kyiv, Ukraine

<sup>3</sup> Universidade Federal de Mato Grosso do Sul, Av. Sen. Felinto. Müller, 1555, C/P. 549, 79074-460, Campo Grande, MS, Brazil

<sup>4</sup> Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, 5001-801, Folhadela, Vila Real, Portugal

<sup>5</sup> Bukovinian State Medical University, 58001, Teatralna Sq., 9, Chernivtsi, Ukraine

<sup>6</sup> Universidade Estadual de Ponta Grossa, Campus de Uvaranas, Av. Gal. Carlos Cavalcanti, 4748, 84030-900, Ponta Grossa, PR, Brazil

<sup>7</sup> Politechnika Lubelska, 20618, Lublin, Poland

<sup>8</sup> Volyn National University, 43000, Voli Ave., 13, Lutsk, Ukraine

<sup>9</sup> Chernivtsi Institute of Trade and Economics of Kyiv National University of Trade and Economics, 58001, Central Sq. 6, Chernivtsi, Ukraine

<sup>10</sup> Faculdade de Engenharia da Universidade do Porto, 4200-465, Rua Dr. Roberto Frias, s/n, Porto, Portugal

<sup>11</sup> Rivne State Basic Medical Academy, 33000, Mykola Karnaukhov Str., 53, Rivne, Ukraine

<sup>12</sup> Abu Ali Ibn Sino Bukhara State Medical Institute, 705018, Navoi Str., 1, Bukhara, Uzbekistan

\* Correspondence: [nightwatcher2401@gmail.com](mailto:nightwatcher2401@gmail.com) (V.V.T.), [ved1988mid@rambler.ru](mailto:ved1988mid@rambler.ru) (P.I.Y.), [zholt1971@ukr.net](mailto:zholt1971@ukr.net) (Z.O.K.);

Scopus Author ID 55758299100

Received: 4.12.2022; Accepted: 30.01.2023; Published: 13.08.2023

**Abstract:** An interesting electrochemical process for sucralose and chloropicrin electrochemical removal has been proposed and theoretically described. The organic halide ion is efficiently removed in mildly acidic media by trivalent vanadium oxyhydroxide, being the nitro group of chloropicrin also thereby reduced. The analysis of the mathematical model confirms that, despite of high probability of the oscillatory behavior, the electrochemical removal of chloropicrin and sucralose is efficient.

**Keywords:** sucralose; chloropicrin; electrochemical removal; vanadium (III) oxyhydroxide; electrochemical oscillations; stable steady-state.


© 2023 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Chloropicrin (Figure 1 to the left side) [1 – 4] is a toxic gas used as an antimicrobial fungicide, herbicide, insecticide, and nematocide in agriculture. It is generally obtained by oxidizing picric acid with hypochlorite in acidic media. Nevertheless, it was also used as a

chemical warfare agent by Germans during World War I and by Russian troops during the Russian-Ukrainian War (2022) [4], which is why it is necessary to rapidly and efficiently remove its remains.

On the other hand, sucralose (Figure 1 to the right) [5 – 9] is an extremely stable artificial sweetener, up to 1000 times as sweet as common sugar. Its effects on human and animal organisms have not been completely studied yet. Moreover, being hardly metabolized, it is accumulated in the environment. Also, the organisms capable of metabolizing it transform the sucralose into toxic chloroorganic derivatives like dioxins and tetrachlorodibenzofurans, which is why the sucralose removal from the environment is also actual.



**Figure 1.** Chloropicrin and sucralose.


Taking into account the presence of organic chlorine in both compounds, the AOP may not be compatible with their removal due to the formation of toxic chlorine gas and chlorine oxides. Cathodic dehalogenation is to be used instead. It leads to the chlorine substitution by hydrogen, leading to the chloride-ion formation [10 – 12]. Also, to impede the chloride ion passage towards the anodic electrolyte, a membrane impeding its permeability is posed to divide the cell into the cathodic and anodic electrolytes. Vanadium (III) oxyhydroxide [13 – 14], acting as a strong reductant in neutral, basic, and mildly acidic media, may be potentially an interesting candidate for cathode modification for sucralose and chloropicrin removal.

Depending on cathode potential and solution pH, the analogous process may also be compatible with the electrochemical reduction of other chloroorganic compounds, like hexachlorobenzene, pentachlorophenol, DDT, dioxins, etc. This is important for environmental remediation, including chemical warfare utilization. The possibility of electroanalytical use of this process, in which the chloroorganic compounds are electrochemically quantified, is also evaluated.

In order to evaluate the process of sucralose and chloropicrin vanadium oxyhydroxide-assisted reduction from the mechanistic point of view, the theoretical description for sucralose and chloropicrin cathodic removal is given in this work. The mathematical model has been developed and analyzed for this system, and the stability requirements and oscillatory and monotonic instability conditions have been established. Also, the behavior of this system has been compared to that of similar systems [15 – 21].

## 2. System and its modeling

Sucralose and chloropicrin are reduced cathodically in acidic media, yielding chloride ions and correspondent hydrocarbon fragments. As for chloropicrin, its nitro group is also reduced, yielding an amino group:



As for vanadium (III) oxyhydroxide, it is regenerated in the electrochemical stage:



Taking this into account, in order to describe the behavior of this system, we introduce three variables:

- c – chloropicrin concentration in the pre-surface layer;
- s – sucralose concentration in the pre-surface layer;
- v – vanadium dioxide surface coverage degree.

Accepting some assumptions [18 – 21], we describe the behavior of the system by equation-set (4):

$$\begin{cases} \frac{dc}{dt} = \frac{2}{\delta} \left( \frac{\Delta}{\delta} (c_0 - c) - r_2 \right) \\ \frac{ds}{dt} = \frac{2}{\delta} \left( \frac{D}{\delta} (s_0 - s) - r_1 \right) \\ \frac{dv}{dt} = \frac{1}{V} (r_1 + r_2 - r_3) \end{cases} \quad (4)$$

Herein  $\Delta$  and  $D$  are diffusion coefficients,  $c_0$  and  $s_0$  are chloropicrin and sucralose bulk concentrations,  $V$  is vanadium dioxide maximal surface concentrations, and the parameters  $r$  are the correspondent reaction rates, calculated as (5 – 6):

$$r_1 = k_1 s \exp(-as) (1 - v)^6 \quad r_2 = k_2 c \exp(-bc) (1 - v)^9 \quad r_3 = k_3 v \exp\left(-\frac{F\varphi_0}{RT}\right) \quad (5 - 7)$$

in which the parameters  $k$  are the correspondent rate constants, parameters  $a$  and  $b$  define the impact of the chemical stages on DEL ionic force,  $F$  is the Faraday number,  $\varphi_0$  stands for zero-charge-related potential slope,  $R$  is the universal gas constant, and  $T$  is the absolute temperature.

Taking into account the chloride-ion formation during the electrochemical process, changing the DEL ionic force, the oscillatory behavior occurs more probably in this system than in similar ones. Nevertheless, the removal process is efficient for converting both compounds, as shown below.

### 3. Results and Discussion

In order to investigate the stability of the system with chloropicrin and sucralose electrochemical conversive removal, we analyze the equation-set (4) by steady-state stability. The steady-state Jacobian matrix members may be described as (8):

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad (8)$$

In which:

$$a_{11} = \frac{2}{\delta} \left( -\frac{\Delta}{\delta} - k_2 \exp(-bc) (1 - v)^9 + bk_2 c \exp(-bc) (1 - v)^9 \right) \quad (9)$$

$$a_{12} = 0 \quad (10)$$

$$a_{13} = \frac{2}{\delta} (9k_2 c \exp(-bc) (1 - v)^8) \quad (11)$$

$$a_{21} = 0 \quad (12)$$

$$a_{22} = \frac{2}{\delta} \left( -\frac{D}{\delta} (s_0 - s) - k_1 \exp(-as) (1 - v)^6 + ak_1 s \exp(-as) (1 - v)^6 \right) \quad (13)$$

$$a_{23} = \frac{2}{\delta} (6k_1s \exp(-as) (1 - v)^5) \quad (14)$$

$$a_{31} = \frac{1}{v} (k_2 \exp(-bc) (1 - v)^9 - bk_2c \exp(-bc)(1 - v)^9) \quad (15)$$

$$a_{32} = \frac{1}{v} (k_1 \exp(-as) (1 - v)^6 - k_1s \exp(-as) (1 - v)^6) \quad (16)$$

$$a_{33} = \frac{1}{v} \left( -6k_1s \exp(-as) (1 - v)^5 - 9k_2c \exp(-bc) (1 - v)^8 - k_3 \exp\left(-\frac{F\varphi_0}{RT}\right) + jk_3v \exp\left(-\frac{F\varphi_0}{RT}\right) \right) \quad (17)$$

Taking into account the main-diagonal elements (9), (13), and (17), we conclude that the oscillatory behavior in this system is possible. Moreover, it becomes more probable than in similar systems [18 – 21] due to the cyclic character of DEL ionic force changes during the chemical and electrochemical stages.

Besides the element  $jk_3v \exp\left(-\frac{F\varphi_0}{RT}\right) > 0$  if  $j>0$ , describing the positive callback by DEL influences of the electrochemical stages, the positive elements responsible for this type of callback are  $bk_2c \exp(-bc) (1 - v)^9 > 0$ , if  $b>0$  and  $ak_1s \exp(-as) (1 - v)^6 > 0$ , if  $a>0$ , both of them referent to the cyclic DEL ionic force, conductivity, and impedance cyclic changes caused by chemical stages. The oscillations amplitude and frequency will be dependent on electrolyte composition. Either way, the oscillations are expected to be relatively frequent and of small amplitude.

Applying the Routh-Hurwitz stability criterion to the equation set (1), we analyze the steady-state stability requirement. In order to avoid cumbersome expressions, we introduce new variables, rewriting thereby the determinant as (18):

$$\frac{4}{\delta^2 v} \begin{vmatrix} -\kappa_1 - \Xi & 0 & \Lambda \\ 0 & -\kappa_2 - \Sigma & P \\ \Xi & \Sigma & -\Lambda - P - \Omega \end{vmatrix} \quad (18)$$

Opening the brackets, applying the Det  $J<0$  requisites, salient from the criterion, and changing the signs to the opposite, we describe the stability requirement as (19):

$$\kappa_1(\kappa_2\Lambda + \kappa_2P + \kappa_2\Omega + \Sigma\Lambda + \Sigma\Omega) + \Xi(\kappa_2\Lambda + \kappa_2P + \Sigma\Omega) > 0 \quad (19)$$

Describing an efficient diffusion-controlled process in which the stable steady-state is easily formed. The inequation (19) is warranted to be satisfied if the kinetical parameters  $\Xi$ ,  $\Sigma$ , and  $\Omega$  are positive, which occurs in most cases. Being those parameters positive, the left side of the inequation (19) gains more positive values. Taking into account that sucralose and chloropicrin do not react with each other and that steady-state stability is easy to obtain and maintain, it is possible to conclude that this process may also serve an electroanalytical purpose, and the steady-state stability will be thereby correspondent to the linear dependence between concentration and current.

As for the monotonic instability, it delimits the margin for steady-state stability and, for electroanalytical effects, the detection limit. Its condition corresponds to the nullity of the determinant, or (20):

$$\kappa_1(\kappa_2\Lambda + \kappa_2P + \kappa_2\Omega + \Sigma\Lambda + \Sigma\Omega) + \Xi(\kappa_2\Lambda + \kappa_2P + \Sigma\Omega) = 0 \quad (20)$$

The model will be accomplished if the chloroorganic compounds react with each other or with a reagent present in the solution. The equation-set (4) will be thereby rewritten as (21):

$$\begin{cases} \frac{dc}{dt} = \frac{2}{\delta} \left( \frac{\Delta}{\delta} (c_0 - c) - r_2 - r_m \right) \\ \frac{ds}{dt} = \frac{2}{\delta} \left( \frac{D}{\delta} (s_0 - s) - r_1 - r_m \right) \\ \frac{dv}{dt} = \frac{1}{V} (r_1 + r_2 - r_3) \end{cases} \quad (21)$$

and its analysis will be analyzed in one of our next works.

#### 4. Conclusions

The system analysis with chloropicrin and sucralose cathodic removal let us conclude that this process is an efficient diffusion-controlled system capable of converting sucralose and chloropicrin into biodegradable compounds economically and green. Furthermore, the easy steady-state stabilization may also have an electroanalytical function, providing facile analytical signal interpretation. As for the oscillatory behavior, it is more probable than for similar systems due to the DEL influences of both chemical and electrochemical processes.

#### Funding

This research received no external funding.

#### Acknowledgments

Volodymyr V. Tkach acknowledges the Engineering Faculty of the University of Porto and the University of Trás-os-Montes and Alto Douro's support during these difficult times for Ukraine and its research.

#### Conflicts of Interest

The authors declare no conflict of interest.

#### References

1. Shi, Zh.; Zhu, J.; Wu, J. *et al.*, Effects of Chloropicrin, Dimethyl Disulfide and Metham Sodium Applied Simultaneously on Soil-Born Bacteria and Fungi. *Agriculture* **2022**, *12*, 1982, <https://doi.org/10.3390/agriculture12121982>.
2. Fang, W.; Wang, Q; Li, Yu. *et al.*, Microbial Regulation of Nitrous Oxide Emissions from Chloropicrin-Fumigated Soil Amended with Biochar. *J. Hazard. Mater.* **2022**, *429*, 128060, <https://doi.org/10.1016/j.jhazmat.2021.128060>.
3. Wang, Y.; Yang, X.; Xu., M. Geissen, V. Effects of Chloropicrin Fumigation and Azoxystrobin Application on Ginger Growth and Phosphorus Uptake. *Ecotoxicol. Env. Safety*, **2022**, *232*, 113246.
4. [www.pravda.com.ua/eng/news/2022/12/4/7379190](http://www.pravda.com.ua/eng/news/2022/12/4/7379190).
5. Debras, Ch; Chazelas, E.; Sellem L. *et al.*, Artificial Sweeteners and Risk of Cardiovascular Diseases: Result from the Prospective NutriNet-Sante Cohort. *BMJ*, **2022**, *378*, e71204, <https://doi.org/10.1136/bmj-2022-071204>.
6. del Pozo, S.; Gómez-Martínez, S.; Díaz, L. *et al.*, Potential Effects of Sucralose and Saccharin on Gut Microbiota: a Review. *Nutrients*, **2022**, *14*, 1682, <https://doi.org/10.3390/nu14081682>.
7. Suez, J.; Cohen, Y.; Váldez-Mas, R. *et al.*, Personalized Microbiome-Driven Effects of Non-Nutritive Sweeteners on Human Glucose Tolerance. *Cell*, **2022**, *185*, 3307 – 3328, <https://doi.org/10.1016/j.cell.2022.07.016>.

8. Méndez-García, L.A.; Bueno-Hernández, N.; Cid-Soto, M.A. *et al.*, Ten-Week Sucralose Consumption Induces Gut Dysbiosis and Altered Glucose and Insulin Levels in Healthy Young Adults. *Microorganisms*, **2022**, *10*, 434, <https://doi.org/10.3390/microorganisms10020434>.
9. Yan, N.; Guo, Zh.; Brusseau, M.L. Sucralose as an Oxidative-Attenuation Tracer for Characterizing the Application of *In Situ* Chemical Oxidation for the Treatment of 1,4-dioxane, *Env. Sci. Proc. Imp.*, **2022**, *8*, 1165 – 1172, <https://doi.org/10.1039/D2EM00185C>.
10. Zhu, X.; Wang, X.; Li, N. *et al.*, Bioelectrochemical System for Dehalogenation: a Review. *Envir. Poll.*, **2022**, *293*, 118519, <https://doi.org/10.1016/j.envpol.2021.118519>.
11. Huang, B.; Sun, Z.; Sun, G. Recent Progress in Cathodic Reduction-Enabled Organic Electrosynthesis: Trends, Challenges and Opportunities. *eScience*, **2022**, *2*, 243 – 277, <https://doi.org/10.1016/j.esci.2022.04.006>.
12. Cantillo, D. Synthesis of Active Pharmaceutical Ingredients Using Electrochemical Methods: Keys to Improve Sustainability. *Chem. Commun.*, **2022**, *58*, 619 – 628, <https://pubs.rsc.org/en/content/articlelanding/2022/cc/d1cc06296d>.
13. Oliveira, R.W.; Gomes, P.; Ribeiro, J.; Reis, J.; Fontenele, M.; Everton, G.; Lyra, W.; Louzeiro, H.; Paula, M.; Filho, V. Determinação espectrofotométrica de Cobre(II) em aguardente de mandioca (Tiquira). *Revista Colombiana de Ciências Químico-Farmacéuticas* **2020**, *49*, 353–371, <https://doi.org/10.15446/rcciquifa.v49n2.89507> (in Portuguese).
14. Bagheri, A.; Hassani Marand, M. Voltammetric and Potentiometric Determination of Cu<sup>2+</sup>, using an Overoxidized Polypyrrole Based Electrochemical Sensor. *Russ. J. Electrochem.* **2020**, *56*, 453–461, <https://doi.org/10.1134/S1023193520060026>.
15. Das, I.; Goel, N.; Agrawal, N.R.; Gupta, S.K. Growth patterns of dendrimers and electric potential oscillations during electropolymerization of pyrrole using mono-and mixed surfactants. *J. Phys. Chem.* **2010**, *114*, 12888–12896, <https://doi.org/10.1021/jp105183q>.
16. Das, I.; Goel, N.; Gupta, S.K.; Agrawal, N.R. Electropolymerization of pyrrole: Dendrimers, nano-sized patterns and oscillations in potential in presence of aromatic and aliphatic surfactants. *Journal of Electroanalytical Chemistry* **2012**, *670*, 1-10, <https://doi.org/10.1016/j.jelechem.2012.01.023>.
17. Bazzouai, M.; Bazzouai, E.A.; Martins, L.; Martins, J.I. Electropolymerization of pyrrole on zinc–lead–silver alloys electrodes in acidic and neutral organic media. *Synthetic Metals* **2002**, *130*, 73-83, [https://doi.org/10.1016/S0379-6779\(02\)00101-7](https://doi.org/10.1016/S0379-6779(02)00101-7).
18. Joshi, N.C.; Malik, S.; Gururani, P. Utilization of Polypyrrole/ZnO Nanocomposite in the Adsorptive Removal of Cu<sup>2+</sup>, Pb<sup>2+</sup> and Cd<sup>2+</sup> Ions from Wastewater. *Letters in Applied NanoBioScience* **2021**, *10*, 2339–2351, <https://doi.org/10.33263/LIANBS103.23392351>.
19. Tkach, V.; Kushnir, M.; de Oliveira, S.; Ivanushko, Y.; Tkach, V.; Mytrofanova, H.; Zadoia, A.; Yagodynets', P.; Kormosh, Z.; Luganska, O. Theoretical Description for an Efficient Rhenium Electrocatalytical Recuperation by Polypyrrole Overoxidation. *Letters in Applied NanoBioScience* **2021**, *10*, 2396-2401, <https://doi.org/10.33263/LIANBS103.23962401>.
20. Tkach, V.V.; Kushnir, M.V.; de Oliveira, S.C.; Parchenko, V.V.; Odyntsova, V.M.; Aksyonova, I.I.; Ivanushko, Y.G.; Yagodynets, P.I.; Kormosh, Z.O. A Descrição Matemática Da Detecção Eletroquímica da Ergina, Assistida Pelos Novos Derivados Triazólicos, Dopados pelo Íon Amavadina. *Rev. Colomb. Cien. Quím. Farm.* **2021**, *50*, 174 – 184, [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0034-74182021000100174&lng=en&nrm=iso](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0034-74182021000100174&lng=en&nrm=iso).
21. Tkach, V.V.; Kushnir, M.V.; Omelchenko, N.M.; Rogozynsky, M.S.; de Oliveira, S.C.; Stotoshchuk, N.M.; Yagodynets, P.I. The Theoretical Description for Amavadin Electrochemical Determination by Poly(Vedelolactone) Doping. *Appl. J. Env. Eng. Sci.* **2019**, *5*, 384 – 389, <https://doi.org/10.48422/IMIST.PRSM/ajeas-v5i4.15374>.